

Intrinsic excitability varies by sex in prepubertal striatal medium spiny neurons

David M. Dorris, Jinyan Cao, Jaime A. Willett, Caitlin A. Hauser and John Meitzen

J Neurophysiol 113:720-729, 2015. First published 5 November 2014; doi:10.1152/jn.00687.2014

You might find this additional info useful...

This article cites 88 articles, 23 of which can be accessed free at:

</content/113/3/720.full.html#ref-list-1>

Updated information and services including high resolution figures, can be found at:

</content/113/3/720.full.html>

Additional material and information about *Journal of Neurophysiology* can be found at:

<http://www.the-aps.org/publications/jn>

This information is current as of February 2, 2015.

Intrinsic excitability varies by sex in prepubertal striatal medium spiny neurons

David M. Dorris,¹ Jinyan Cao,^{1,2} Jaime A. Willett,^{1,2,3} Caitlin A. Hauser,¹ and John Meitzen^{1,2,4,5}

¹Department of Biological Sciences, North Carolina State University, Raleigh, North Carolina; ²W. M. Keck Center for Behavioral Biology, North Carolina State University, Raleigh, North Carolina; ³Graduate Program in Physiology, North Carolina State University, Raleigh, North Carolina; ⁴Center for Human Health and the Environment, Center for Comparative Medicine and Translational Research, North Carolina State University, Raleigh, North Carolina; and ⁵Grass Laboratory, Marine Biological Laboratory, Woods Hole, Massachusetts

Submitted 12 September 2014; accepted in final form 3 November 2014

Dorris DM, Cao J, Willett JA, Hauser CA, Meitzen J. Intrinsic excitability varies by sex in prepubertal striatal medium spiny neurons. *J Neurophysiol* 113: 720–729, 2015. First published November 5, 2014; doi:10.1152/jn.00687.2014.—Sex differences in neuron electrophysiological properties were traditionally associated with brain regions directly involved in reproduction in adult, postpubertal animals. There is growing acknowledgement that sex differences can exist in other developmental periods and brain regions as well. This includes the dorsal striatum (caudate/putamen), which shows robust sex differences in gene expression, neuromodulator action (including dopamine and 17 β -estradiol), and relevant sensorimotor behaviors and pathologies such as the responsiveness to drugs of abuse. Here we examine whether these sex differences extend to striatal neuron electrophysiology. We test the hypothesis that passive and active medium spiny neuron (MSN) electrophysiological properties in prepubertal rat dorsal striatum differ by sex. We made whole cell recordings from male and female MSNs from acute brain slices. The slope of the evoked firing rate to current injection curve was increased in MSNs recorded from females compared with males. The initial action potential firing rate was increased in MSNs recorded from females compared with males. Action potential after-hyperpolarization peak was decreased, and threshold was hyperpolarized in MSNs recorded from females compared with males. No sex differences in passive electrophysiological properties or miniature excitatory synaptic currents were detected. These findings indicate that MSN excitability is increased in prepubertal females compared with males, providing a new mechanism that potentially contributes to generating sex differences in striatal-mediated processes. Broadly, these findings demonstrate that sex differences in neuron electrophysiological properties can exist prepuberty in brain regions not directly related to reproduction.

intrinsic excitability; sex differences; medium spiny neuron; dorsal striatum; mEPSC

NEURAL SEX DIFFERENCES ARE well established in many vertebrate brain regions, especially those directly involved in reproduction in adult, postpubertal animals (Breedlove and Hampson 2002; De Vries 2004; Yang and Shah 2014). Examples of these include the sexually dimorphic nucleus of the preoptic area (SDN) (Gorski et al. 1978), the spinal nucleus of the bulbocavernosus (SNB) (Breedlove and Arnold 1981), and the telencephalic song control nuclei in sexually dimorphic songbirds (Nottebohm and Arnold 1976). These now famous brain regions all show robust sex differences in neuroanatomy and

physiology and clear behavioral relevance to sex specific behaviors. What is less clear is the extent of sex differences in basic neurophysiological properties in brain regions without such dramatic sex differences in neuroanatomy, especially during the prepubertal period widely used for electrophysiological recordings. This is an important and timely question, given the current debate regarding how to properly account for sex in both basic and clinical biomedical science (Arnold and Lusis 2012; Beery and Zucker 2011; Cahill 2006; Geller et al. 2011; Prendergast et al. 2014; Woodruff et al. 2014), the known sex differences in many neural pathologies (Becker et al. 2013; Cosgrove et al. 2007; Giorgi et al. 2014; Ober et al. 2008), and the growing literature for sex differences in synaptic organization/neuromodulation across the nervous system (Babayán and Kramar 2013; Cooke and Woolley 2005; Huang and Woolley 2012; Mermelstein et al. 1996; Nunez and McCarthy 2008; Remage-Healey 2014; Srivastava et al. 2010).

We thus chose to investigate sex differences in the rat dorsal striatum (caudate/putamen). This brain region was targeted for both its prominence and the known sex differences in striatal-mediated behaviors and pathologies. These include significant sex differences in steroid sex hormone influences on sensorimotor function and behaviors, impulsivity, and striatal-mediated learning (Becker 2002; Calhoun 1962; Eckel et al. 2000; Hosseini-Kamkar and Morton 2014; Zurkovsky et al. 2007). Regarding pathologies, robust sex differences exist in the responsiveness to drugs of abuse in both humans and rats (Becker and Hu 2008; Becker et al. 2013; Bobzean et al. 2014; Carroll and Anker 2010; Fattore et al. 2014). Across models, females exhibit increased locomotor sensitivity, escalation, and motivation to take psychostimulants after initial exposure compared with males, with estradiol playing a significant mechanistic role. Interestingly, the rat dorsal striatum and nucleus accumbens express little to no nuclear estrogen receptors, and instead express membrane-associated estrogen receptors α and β and GPER-1 (Almey et al. 2012; Grove-Strawser et al. 2010; Kuppens and Beyer 1999; Mermelstein et al. 1996; Schultz et al. 2009).

Given the importance of these behaviors and pathologies, much research has established sex differences in adult striatum gene expression (Chen et al. 2009; Ghahramani et al. 2014; Trabzuni et al. 2013), estradiol sensitivity (Cummings et al. 2014; Grove-Strawser et al. 2010; Mermelstein et al. 1996; Schultz et al. 2009), catecholamine action (Becker and Hu 2008; Becker et al. 2013; Di Paolo 1994; Meitzen et al. 2013),

Address for reprint requests and other correspondence: J. Meitzen, Dept. of Biological Sciences, NC State Univ., Campus Box 7617, Raleigh, NC 27695-7617 (e-mail: jemeitze@ncsu.edu).

Δ FosB expression (Sato et al. 2011), and GABA and dopamine release (Becker 1990; Hu et al. 2006; Walker et al. 2000; Xiao and Becker 1998). In a related brain region, the nucleus accumbens, sex differences in synaptic organization have been found (Forlano and Woolley 2010; Wissman et al. 2012; Wissman et al. 2011). In contrast to regions like the SDN, SNB, and sexually dimorphic song control nuclei, striatal brain regions show no sex differences in neuron density or soma size (Meitzen et al. 2011), and the volume of the nucleus accumbens does not differ by sex (Campi et al. 2013). Most notably, it is unknown whether the basic electrophysiological properties of striatal neurons differ by sex.

Here we test the hypothesis that passive and active medium spiny neuron (MSN) electrophysiological properties in prepubertal rat dorsal striatum (caudate/putamen) differ by sex. The prepubertal period was chosen as it is widely used for electrophysiological recordings. We raised male and female rats and then recorded from MSNs using whole cell current clamp configuration in acute brain slices of dorsal striatum. We found that the active electrophysiological properties varied by sex, with female MSNs exhibiting increased intrinsic excitability compared with male MSNs. No sex differences in passive electrophysiological properties or miniature excitatory synaptic currents (mEPSCs) were detected. These findings provide insight on a new mechanism that potentially contributes to sex differences in striatal-mediated behavior and pathologies. Broadly, these findings demonstrate that sex differences in neuron electrophysiological properties can exist prepuberty in brain regions not directly related to reproduction.

MATERIALS AND METHODS

Animals

All animal protocols were approved by Institutional Animal Care and Use Committee at North Carolina State University or the Marine Biological Laboratory. Experiments took place at both institutions. Female ($n = 18$) and male ($n = 18$) Sprague-Dawley CD IGS rats were born from timed-pregnant females purchased from Charles River (Raleigh, NC). Rats were housed with their littermates and dam until weaning on postnatal day 21 (P21), and afterward with same-sex littermates. Age at experimental use ranged from P11 to P23 and was matched between sexes (mean \pm SE: male, $P15 \pm 1$; female, $P15 \pm 1$). All cages were washed polysulfone (BPA-free) and were filled with bedding manufactured from virgin hardwood chips (Beta Chip; NEPCO, Warrensburg, NY) to avoid the endocrine disruptors present in corn cob bedding (Mani et al. 2005; Markaverich et al. 2002; Villalon Landeros et al. 2012). Rooms were temperature, humidity, and light controlled (23°C, 40% humidity, 12 h light-12 h darkness cycle). Soy protein-free rodent chow (2020X; Teklad, Madison, WI) and glass-bottle provided water were available ad libitum.

Electrophysiology

Preparation of brain slices. Methods for preparing brain slices for electrophysiological recordings were as previously described (Dorris et al. 2014). Rats were deeply anesthetized with isoflurane gas and killed by decapitation. The brain was dissected rapidly into ice-cold, oxygenated sucrose artificial cerebrospinal fluid (ACSF) containing (in mM): 75 sucrose, 1.25 NaH_2PO_4 , 3 MgCl_2 , 0.5 CaCl_2 , 2.4 Na pyruvate, 1.3 ascorbic acid from Sigma-Aldrich (St. Louis, MO), and 75 NaCl, 25 NaHCO_3 , 15 dextrose, 2 KCl from Fisher (Pittsburg, PA); osmolarity 295–305 mOsm, pH 7.2–7.4. Coronal brain slices (300 μm) were prepared using a vibratome and then incubated in

regular ACSF containing (in mM): 126 NaCl, 26 NaHCO_3 , 10 dextrose, 3 KCl, 1.25 NaH_2PO_4 , 1 MgCl_2 , 2 CaCl_2 , 295–305 mOsm, pH 7.2–7.4, for 30 min at 35°C, and at least 30 min at room temperature (21–23°C). Slices were stored submerged in room temperature, oxygenated ACSF for up to 5 h after sectioning in a large volume bath holder.

Electrophysiological recording. After resting for ≥ 1 h after sectioning, slices were placed in a Zeiss Axioscope equipped with IR-DIC optics, a Dage IR-1000 video camera, and $\times 10$ and $\times 40$ lenses with optical zoom. Slices were superfused with oxygenated ACSF heated to 27 ± 1 °C (male 27 ± 1 °C, female 27 ± 1 °C; $P > 0.05$). In some experiments, ACSF contained the GABA_A receptor antagonist picrotoxin (PTX, 150 μM ; Fisher), the NMDA receptor antagonist D-AP5 (10 μM , Sigma-Aldrich), and the AMPA receptor antagonist DNQX (25 μM ; Tocris, Minneapolis, MN). Whole cell patch-clamp recordings were made from MSNs in the dorsal striatum via glass electrodes (4–8 M Ω) containing (in mM): 115 K D-gluconate, 8 NaCl, 2 EGTA, 2 MgCl_2 , 2 MgATP, 0.3 NaGTP, 10 phosphocreatine from Sigma-Aldrich and 10 HEPES from Fisher, 285 mOsm, pH 7.2–7.4. Signals were amplified, filtered (2 kHz), and digitized (10 kHz) with a MultiClamp 700B amplifier attached to a Digidata 1550 system and a personal computer using pClamp 10 software. Membrane potentials were corrected for a calculated liquid junction potential of -13.5 mV. Using previously described procedures (Farries et al. 2005; Meitzen et al. 2009), recordings were made in current clamp to assess neuronal electrophysiological properties. MSNs were identified by their medium-sized somas, the presence of a slow ramping subthreshold depolarization in response to low-magnitude positive current injections, a hyperpolarized resting potential more negative than -65 mV, inward rectification, and prominent spike after-hyperpolarization (AHP) (Belleau and Warren 2000; O'Donnell and Grace 1993).

In a subset of recordings, MSNs were then voltage-clamped at -70 mV and mEPSCs recorded in the presence of tetrodotoxin (TTX, 1 μM ; Abcam Biochemicals) and PTX (150 μM). These recording parameters make the recorded mEPSCs to be most likely AMPA receptor mediated. mEPSCs were recorded for at least 5 min, with the exception of one female neuron that was recorded for 3 min. Input and series resistance was monitored for possible changes, and cells were discarded if input or series resistance changed $>15\%$.

Data Analysis

Basic electrophysiological properties and action potential (AP) characteristics were analyzed with pClamp 10. After break-in, the resting membrane potential was first allowed to stabilize ~ 1 –2 min, as in (Mu et al. 2010). We then injected at least three series of depolarizing and hyperpolarizing current injections to elicit basic neurophysiological properties (Meitzen et al. 2009). For most properties measured, we followed the definitions of (Meitzen et al. 2009), which were drawn from those of Farries and colleagues (Farries et al. 2005; Farries and Perkel 2000; 2002). Following Farries, for each neuron, measurements were made of at least five APs generated from minimal current injections. These measurements were then averaged to generate the reported AP measurement for that neuron. For AP measurements, only the first generated AP was used unless more APs were required to meet the standard five APs per neuron. We used different methods from Farries to calculate AP threshold, steady-state firing rate, rectified range input resistance, inward rectification, and percent inward rectification. AP threshold was defined as the first point of sustained positive acceleration of voltage ($\delta^2V/\delta t^2$) that was also more than three times the SD of membrane noise before the detected threshold (Baufreton et al. 2005). We defined steady-state firing rate as the mean firing rate over the last 300 ms of the current pulse (Gale and Perkel 2006). Initial firing rate was defined as the inverse of the first interspike interval. Rectified range input resistance, inward rectification, and percent inward rectification were calculated as described previously (Belleau and Warren 2000). The slope of the

evoked firing rate to positive current curve (FI slope) was calculated from the first current that evoked an AP to the first current that generated the maximum evoked firing rate (Meitzen et al. 2009). Input resistance in the linear, nonrectified range was calculated from the steady-state membrane potential in response to -0.02 nA hyperpolarizing pulses. The membrane time constant was calculated by fitting a single exponential curve to the membrane potential change in response to -0.02 nA hyperpolarizing pulses. Membrane capacitance was calculated using the following equation: capacitance = membrane time constant/input resistance. Sag index was used to assess possible sex differences in hyperpolarization-induced "sag" (i.e., I_H current) (Farries et al. 2005). Sag index is the difference between the minimum voltage measured during the largest hyperpolarizing current pulse and the steady-state voltage deflection of that pulse, divided by the steady-state voltage deflection. Thus, a cell with no sag would have a sag index of 0, whereas a cell whose maximum voltage deflection is twice that of the steady-state deflection would have a sag index of 1. Cells with considerable sag typically have an index of ≥ 0.1 .

mEPSC frequency, amplitude, and decay were analyzed off-line with Mini Analysis (Synaptosoft, <http://www.synaptosoft.com/MiniAnalysis/>). Threshold was set at 2.5, the value of the root mean square of 10 blocks of the baseline noise with a minimum value of 5 pA, and accurate event detection was validated by visual inspection. There were no differences in root mean square noise between sexes (male 1.1 ± 0.01 , female 1.5 ± 0.02 ; $t_{(19)} = 1.923$; $P > 0.05$).

Statistics

Experiments were analyzed by two-tailed t -tests, Mann-Whitney tests, Kolmogorov-Smirnov tests, one- or two-way ANOVAs, linear regressions, or ANCOVAs (Excel 2010, Microsoft, Redmond, WA, or Prism version 5.00, GraphPad Software, La Jolla, CA). Distributions were analyzed for normality by the D'Agostino and Pearson omnibus normality test, and t -tests or Mann-Whitney tests were employed as appropriate. The use of t -tests or Mann-Whitney tests did not alter overall experimental conclusions. P values < 0.05 were considered a priori as significant. Data are presented as means \pm SE.

RESULTS

We recorded from 33 MSNs from 18 prepubertal male rats and 32 MSNs from 18 prepubertal female rats. MSNs are the predominant neuron type in the dorsal striatum, projecting both within and outside the brain region. MSN electrophysiological properties closely resembled those reported in earlier studies that used males or animals of undetermined sex, including the presence of a slow ramping subthreshold depolarization in response to low-magnitude positive current injections, a hyperpolarized resting potential, inward rectification, and prominent spike AHP (Fig. 1A, Table 1) (Belleau and Warren 2000; Farries et al. 2005; O'Donnell and Grace 1993; Shen et al. 2004).

Female MSNs Show Increased Evoked Firing Rates Compared With Male Neurons

We then tested the hypothesis that MSN electrophysiological properties varied between males and females by injecting MSNs with a series of positive and negative currents and assessing standard electrophysiological properties (Table 1). Several electrophysiological properties related to intrinsic excitability varied by sex, collectively indicating that female MSNs showed increased excitability compared with male MSNs. Action potential firing rates evoked by depolarizing

current injection were visibly increased in MSNs recorded from female rats compared with male rats (Fig. 1, A and B). We quantified this by comparing the slope of the evoked firing rate to positive current curve (FI slope) between male and female MSNs (Fig. 1C). MSNs from female animals showed a steeper FI slope compared with MSNs recorded from male animals ($t_{(61)} = 3.026$, $P < 0.004$). This indicates that female neurons fire more APs per ampere of injected current throughout the linear range, reaching maximum rates more quickly. Supporting this finding, the cumulative distribution of female MSN FI slopes also differed from that recorded from male MSNs (Fig. 1D; $D_{(61)} = 0.3909$, $P < 0.02$). These data indicate that MSNs show increased excitability in females compared with males.

There was no difference in maximum firing rate between MSNs recorded from males and females (male 16.7 ± 1.7 Hz, female 15.2 ± 1.2 Hz; $t_{(61)} = 0.70$, $P > 0.05$), suggesting that the sex difference in AP generation likely involves either the delay to first spike or the initial firing rate (initial: firing rate of the first interspike interval, early in the current injection). No sex differences were detected in the delay to first spike at either the minimum current injection necessary for AP generation (Table 1), or at $\Delta 10$ pA from the minimum current injection necessary for AP generation (male 99 ± 7 ms, female 99 ± 9 ; $P > 0.05$). Instead, female MSNs showed increased initial firing rates compared with male MSNs (Fig. 1E; sex $F_{(1, 259)} = 7.014$, $P < 0.009$; current $F_{(5, 259)} = 24.22$, $P < 0.0001$; interaction $F_{(5, 259)} = 0.1556$, $P > 0.05$). We note that one outlier ($> +4$ SDs from the mean) in the female group was not included in analysis of the instantaneous firing rate. Removal of this outlier did not change experimental conclusions (statistics including outlier: sex $F_{(1, 265)} = 9.251$, $P < 0.003$; current $F_{(5, 265)} = 3.687$, $P < 0.004$; interaction $F_{(5, 265)} = 0.0852$, $P > 0.05$). Conversely, while the overall magnitude of the steady state firing rate (steady state: mean firing rate over the last 300 ms of the current injection) was increased in female MSNs compared with male MSNs across current injections, this did not reach statistical significance (Fig. 1F; sex $F_{(1, 265)} = 2.616$, $P > 0.05$; current $F_{(5, 265)} = 8.187$, $P < 0.0001$; interaction $F_{(5, 265)} = 0.1030$, $P > 0.05$).

Female MSNs Show Decreased AHP and Hyperpolarized AP Threshold Compared With Male Neurons

A sex difference in evoked firing rate suggests that there may be differences in either AP properties or the passive membrane properties of the MSN. Regarding AP properties, female MSNs showed decreased magnitude of the AHP peak compared with male MSNs (Fig. 1G, $t_{(61)} = 3.035$, $P < 0.004$). The cumulative distribution of AHP peaks recorded from female MSNs also differed from those recorded from male MSNs (Fig. 1H, $D_{(61)} = 0.3212$, $P < 0.005$). The AP threshold also differed by sex, with female MSNs showing a hyperpolarized AP threshold compared with male MSNs (Fig. 1I, $t_{(61)} = 2.099$; $P < 0.041$). The cumulative distribution of female MSN AP thresholds also differed from those recorded from male MSNs (Fig. 1J, $D_{(61)} = 0.2697$, $P < 0.05$). No other AP property differed by sex (Table 1). These data support the hypothesis that decreased AHP peak and hyperpolarized AP potentials comprise part of the mechanism underlying increased excitability in female MSNs, especially given that both




Fig. 1. Evoked firing rate is increased in medium spiny neurons (MSNs) recorded from prepubertal rat females compared with males in dorsal striatum, and this is driven by sex differences in action potential properties. *A*: response of a MSN from a male (*left*) and from a female (*right*) to depolarizing current injections. *B*: evoked firing rate and injected current (FI) curves of individual MSNs differ by sex. *C*: MSNs recorded from females show an increased slope of the evoked firing rate to injected current curve (FI slope). *D*: the cumulative frequency distribution of MSN FI slopes is shifted to the right in females. *E*: mean initial firing rate (the reciprocal of the 1st interspike interval) is increased in MSNs recorded from females. *F*: mean steady-state firing rate (mean firing rate over the last 300 ms of the current injection) does not significantly differ by sex. However, we do note that the mean steady-state firing rate is consistently higher in females than in males. *G*: action potential (AP) after-hyperpolarization (AHP) peak amplitude is decreased in females. *H*: the cumulative frequency distribution of MSN AHP peak amplitude is shifted to the right in females. *I*: AP threshold is hyperpolarized in females. *J*: the cumulative frequency distribution of MSN AP threshold is shifted to the left in females. The P value within each subpanel indicates statistical significance; complete statistical information is in RESULTS.

of these properties are associated with changes in MSN excitability in other contexts (Mu et al. 2010; Shen et al. 2005).

We reasoned that if these sex differences in AP properties contribute to sex differences in FI slope, then these properties should be correlated. We thus calculated linear regressions between MSN FI slopes and, respectively, AHP peaks and AP thresholds. Increased FI slopes strongly associated with decreased AHP peaks (Fig. 2*A*; slope 0.02, $r^2 = 0.35$, $P < 0.001$). Likewise, FI slope and AP threshold also correlated, with increased FI slopes associating with hyperpolarized AP thresholds (Fig. 2*B*; slope -0.02 , $r^2 = 0.08$, $P < 0.03$). To validate this methodology, we also calculated a linear regression between AHP peak and AP threshold, finding that decreased AHP peaks associated with hyperpolarized thresholds (Fig. 2*C*, slope -1.22 , $r^2 = 0.35$, $P < 0.0001$). This makes sense given

that female MSNs exhibit both of these properties and increased FI slope. These data indicate that decreased AHP peak and hyperpolarized AP threshold are associated with increased neuron excitability.

In addition to AP properties, other mechanisms that could potentially drive differences in MSN excitability are changes in passive membrane properties such as the input resistance or the membrane time constant. Passive membrane properties were not different between male and female MSNs (Table 1). This included input resistance in both the linear and rectified ranges (Fig. 3; $F_{(3,570)} = 0.02657$, $P > 0.05$), the membrane time constant, and capacitance (Table 1). The lack of sex differences in capacitance is consistent with a previous report showing no sex difference in MSN soma size in rat dorsal striatum (Meitzen et al. 2011). These results support the hypothesis that sex

Table 1. *Electrophysiological properties of male and female medium spiny neurons*

Property	Male	Female	Statistics (<i>t/U</i> , <i>P</i>)
Resting potential, mV	-81.2 ± 0.9 (33)	-80.0 ± 1.2 (31)	0.52, 0.40
Input resistance, $M\Omega$	448 ± 42 (33)	465 ± 42 (31)	484, 0.72
Time constant of the membrane, ms	23 ± 1 (33)	27 ± 2 (31)	1.74, 0.08
Capacitance, pF	61 ± 4 (33)	69 ± 7 (31)	476, 0.64
Rectified range input Resistance, $M\Omega$	289 ± 29 (33)	291 ± 26 (31)	499, 0.87
Inward rectification, $M\Omega$	156 ± 18 (33)	175 ± 19 (31)	0.72, 0.47
% Inward rectification, %	67 ± 2 (33)	65 ± 2 (31)	0.86, 0.39
Sag index	0.01 ± 0.00 (33)	0.01 ± 0.00 (31)	501, 0.89
AP threshold, mV	-54 ± 1 (33)	-58 ± 1 (30)	2.09, 0.040
AP amplitude, mV	72 ± 2 (33)	75 ± 2 (30)	1.26, 0.21
AP width at half-peak, ms	2.6 ± 0.1 (33)	2.6 ± 0.1 (30)	0.27, 0.78
AHP peak, mV	-10.8 ± 0.7 (33)	-8.3 ± 0.4 (30)	3.04, 0.004
AHP time to peak, ms	52.9 ± 2.9 (33)	47.9 ± 4.1 (30)	1.18, 0.24
Delay to first spike, ms	313 ± 15 (33)	307 ± 19 (30)	0.24, 0.81
Rheobase, nA	0.040 ± 0.008 (33)	0.025 ± 0.004 (30)	356, 0.053
FI slope, Hz/nA	312.1 ± 18.5 (33)	395.2 ± 20.4 (30)	3.03, 0.004

Values are means \pm SE. Numbers in parentheses indicate sample size. The sag index is unitless. None of these neurons fired spontaneous action potentials. Statistical differences between groups are depicted in boldface. Data were analyzed with *t*-tests or Mann-Whitney *U*-tests as appropriate. AP, action potential; AHP, afterhyperpolarization; FI, frequency of evoked spikes to injected depolarization current.

differences in MSN excitability are driven by differences in action potential properties and not passive membrane properties.

Sex Differences Are Not Blocked by Glutamate and GABA Receptor Antagonists

Our working model regarding these particular sex differences in MSN electrophysiological properties is that they are intrinsic to the neuron and not driven by external synaptic input. If this interpretation is correct, then sex differences should be preserved during blockade of glutamatergic and GABAergic receptors. To test this hypothesis, we exposed a subset of MSNs to a cocktail of the NMDA receptor antagonist D-AP5 (10 μ M), the AMPA receptor antagonist DNQX (25 μ M), and the GABA_A receptor antagonist PTX (150 μ M). This drug combination eliminated spontaneous postsynaptic potentials (Fig. 4A), indicating effective blockade of glutamatergic and GABAergic receptors. We then assessed standard electrophysiological properties as described above. Exposure to D-AP5, DNQX, and PTX did not eliminate sex differences in either FI slope (Fig. 4B; sex $F_{(1, 8)} = 13.92$, $P < 0.03$; drug $F_{(2, 8)} = 1.78$, $P > 0.05$), AHP peak (Fig. 4C; sex $F_{(1, 8)} = 13.92$, $P < 0.045$; drug $F_{(2, 8)} = 1.78$, $P > 0.05$) or AP threshold (Fig. 4D; $F_{(1, 8)} = 13.92$, $P < 0.02$; drug $F_{(2, 8)} = 0.07$, $P > 0.05$). Exposure to D-AP5, DNQX, and PTX did not alter any other measured electrophysiological property (data not shown). We also note that the sampled male neurons by chance exhibited decreased FI slopes and increased magnitudes of the AHP peak compared with the overall data set (analysis




Fig. 2. Sex differences in AP properties correlate with sex differences in FI slope. *A*: decreased AHP peaks associate with increased FI slopes. *B*: hyperpolarized AP thresholds associate with increased FI slopes. *C*: hyperpolarized AP thresholds associate with decreased AHP peaks. The *P* value within each panel indicates statistical significance; complete statistical information is in Results.

not shown). Overall, these data support the hypothesis that sex differences in MSN electrophysiological properties are intrinsic and not driven by glutamatergic or GABAergic synaptic input.




Fig. 3. No sex difference is detected in input resistance in either the linear or rectified range. See RESULTS and Table 1 for statistical analysis.




Fig. 4. Sex differences are not driven by synaptic input or by age at recording. *A*: example blockade of miniature postsynaptic potentials via simultaneous application of picrotoxin (PTX), DNQX, and AP-5 to block GABAergic and glutamatergic synaptic activity. *B*: sex differences in FI slope were not blocked by drug application. *C*: sex differences in AHP peak were not blocked by drug application. *D*: sex differences in AP threshold were not blocked by drug application. *E*: relationship of FI slope and age of the animal at recording. The linear regression of the female data set shows an increased intercept compared with the male. *F*: relationship between AHP peak and age. The linear regression of the female data set shows a different slope compared with the male. *G*: relationship of AP threshold and age. The linear regression of the female data set shows a hyperpolarized intercept compared with the male. The P value within each panel indicates statistical significance; complete statistical information is in RESULTS.

Sex Differences Are Present Across All Examined Ages

A priori, we hypothesized that sex differences in electrophysiological properties would be stable throughout the recorded prepubertal period. This hypothesis was chosen given that MSNs reach electrophysiological maturity after the initial critical period for organizational steroid sex hormone action, which may program sex differences in the dorsal striatum and relevant behaviors/pathologies. To test the interaction of sex differences in MSN electrophysiological properties with age at recording, MSNs were analyzed across a sex-matched age range that encompassed the late prepubertal period (male $P15 \pm 1$, female $P15 \pm 1$; $t_{(65)} = 0.36$, $P > 0.05$). The age range began with the onset of the presence of mature MSNs ($\sim P11$) to prior to weaning ($\sim P20$) to just before the beginning of the peripubertal period ($\sim P22$ – $P24$). We then calculated linear regressions between age and male and female MSN FI slope, AHP peak, and AP threshold. We first analyzed whether the slopes of the linear regressions were significantly nonzero to determine whether the measured property differed by age. We then analyzed whether the slopes and elevations/intercepts differed by sex.

FI slope decreased with age in both males and females (Fig. 4*E*) (male: slope -11.82 , $r^2 = 0.14$, $P < 0.05$; female: slope -12.24 , $r^2 = 0.13$, $P < 0.05$), similar to previously reported results in MSNs recorded from rat nucleus accumbens of unknown sex (Belleau and Warren 2000). The slopes of the linear regressions did not differ by sex ($F_{(1,59)} = 0.00$, $P > 0.05$). However, the intercept of the regression did vary by sex, with females showing increased FI slope across all ages (male intercept 496, female intercept 571; $F_{(1,60)} = 6.84$, $P < 0.02$). AHP peak amplitude remained stable with age in females but not in males (Fig. 4*F*; male slope -0.453 , $r^2 = 0.14$, $P < 0.03$; female slope 0.04, $r^2 = 0.00$, $P > 0.05$). Thus, the slopes of the linear regressions differed by sex ($F_{(1,59)} = 4.14$, $P < 0.05$). AP threshold did not change across age in either males or females (Fig. 4*G*, male slope -0.06 , $r^2 = 0.00$, $P > 0.05$; female slope -0.36 , $r^2 = 0.03$, $P > 0.05$), similar to previously reported results of accumbal and dorsal striatal MSNs of unknown sex (Belleau and Warren 2000; Tepper et al. 1998). Accordingly, the slopes of the linear regressions did not differ by sex ($F_{(1,59)} = 0.30$, $P > 0.05$). The intercept did differ by sex, with females showing hyperpolarized AP threshold across all ages compared with males (male intercept -53.08 , female




Fig. 5. No sex differences are detected in miniature excitatory synaptic current (mEPSC) properties. *A*: representative examples of mEPSCs recorded in male and female dorsal striatum MSNs. MSNs were voltage clamped at -70 mV and recorded in the presence of tetrodotoxin (TTX) and PTX to block voltage-gated sodium channels and GABAergic synaptic activity, respectively. *B, left*: no sex difference was detected in MSN mEPSC frequencies. *Right*: no sex differences were detected in the cumulative frequency distribution of MSN mEPSC frequencies. *C, left*: no sex difference was detected in MSN mEPSC amplitude. *Right*: no sex difference was detected in the cumulative frequency distribution of MSN mEPSC amplitudes. *D, left*: no sex difference was detected in MSN mEPSC decay time. *Right*: no sex difference was detected in the cumulative frequency distribution of MSN mEPSC decay times. The P value within each subpanel indicates statistical significance; complete statistical information is in RESULTS.

intercept -52.62 ; $F_{(1,60)} = 4.78$, $P < 0.035$). We note that sex differences in the linear regressions of FI slope, AHP peak, and AP threshold are maintained when postweaning animals are excluded from analysis (data not shown). Collectively, these results indicate that sex differences in MSN electrophysiological properties are determined early in development and are already present in the prepubertal period analyzed here.

No Sex Differences Are Detected in mEPSC Properties

In a subset of recordings, we voltage-clamped 10 male and 11 female MSNs to -70 mV and recorded mEPSCs in the presence of TTX and PTX (Fig. 5*A*). We then analyzed mEPSC frequency, amplitude, and decay (Table 2). This was spurred by the recognition that intrinsic electrophysiological properties act in concert with synaptic properties to generate neuronal output and by a previous report of sex differences in mEPSC properties in adult rat nucleus accumbens core (Wissman et al. 2011). No sex differences were detected in mEPSC frequency (Fig. 5*B*; $t_{(19)} = 1.632$, $P > 0.05$), mEPSC amplitude (Fig. 5*C*; $t_{(19)} = 0.5259$, $P > 0.05$), or mEPSC decay (Fig. 5*D*; $t_{(19)} = 1.078$, $P > 0.05$). To further test this conclusion, we then analyzed mEPSC properties by age at recording. To do this, we calculated linear regressions between age and male and female MSN mEPSC frequency, amplitude, and decay. We first analyzed whether the slopes of the linear regressions were significantly nonzero to determine whether the measured property differed by age. We found no slopes that were significantly nonzero, indicating little change over the analyzed age range (data not shown). We then analyzed whether the slopes and elevations/intercepts of mEPSC frequency, amplitude, and decay differed by sex. No differences were detected (data not shown). Overall, these data indicate that sex differences in prepubertal MSN do not include mEPSC properties.

DISCUSSION

There are five general findings of these experiments. First, the slope of the evoked firing rate to current injection curve was increased in MSNs recorded from females compared with males. Second, the initial action potential firing rate was increased in MSNs recorded from females. Third, AP AHP peak was decreased and AP threshold hyperpolarized in MSNs recorded from females. Fourth, no sex differences in passive electrophysiological or mEPSC properties were detected. Fifth, these sex differences were not attenuated by GABAergic or glutamatergic receptor blockade and were present across all ages examined. These findings indicate that MSN intrinsic membrane excitability in the dorsal striatum is increased in prepubertal females compared with males. Broadly, these find-

Table 2. mEPSC properties of male and female medium spiny neurons

mEPSC Property	Male	Female	Statistics (t , P)
Frequency, Hz	1.7 ± 0.4 (10)	3.1 ± 0.9 (11)	1.36, 0.19
Amplitude, pA	9.5 ± 0.6 (10)	9.5 ± 0.6 (11)	0.53, 0.61
Decay, ms	5.9 ± 0.7 (10)	4.9 ± 0.7 (11)	1.08, 0.29

Values are means \pm SE. Numbers in parentheses indicate sample size. No significant differences were detected. mEPSC, miniature excitatory synaptic current.

ings show that sex differences in neuron electrophysiological properties can occur in brain regions not directly related to reproduction and during the prepubertal developmental period in which many neuron electrophysiological recordings take place.

To our knowledge this is the first report of a sex difference in an intrinsic electrophysiological property of the MSN, providing a new potential mechanism for the known sex differences in striatal-mediated behavior and pathologies. Intrinsic membrane excitability regulates AP generation in response to synaptic input (Hille 2001), making intrinsic excitability a key player in determining the functional output of striatal circuitry and a final common convergence point of all striatal sex differences. Multiple ion channels have been implicated in generating differences in intrinsic excitability, with particular targets in MSNs including soma-expressed sodium channels, calcium-activated potassium channels, and L-type calcium channels (Hille 2001; Kole and Stuart 2012; Kourrich and Thomas 2009; Mermelstein et al. 1996; Mu et al. 2010; Zhang et al. 1998). Determining which of these drive sex differences in intrinsic excitability is an important future extension of this study.

Intrinsic excitability is strongly implicated in both normal and pathological striatal function (Ishikawa et al. 2009; Kourrich and Thomas 2009; Mu et al. 2010; Wolf 2010; Zhang et al. 1998). Indeed, previous research has not only found differences in MSN intrinsic membrane excitability related to homeostatic plasticity (Ishikawa et al. 2009) and drugs of abuse (Kourrich and Thomas 2009; Mu et al. 2010; Wolf 2010; Zhang et al. 1998), but also differences in intrinsic excitability between D1- and D2-dopamine receptor expressing MSNs (Gertler et al. 2008; Planert et al. 2013). These differences in intrinsic excitability between MSN subtypes are already present in rats during the prepubertal period examined here. One possibility is that sex differences in MSN intrinsic excitability were generated by differential sampling of D1- or D2-dopamine receptor expressing MSNs. We believe this to be unlikely. In rats of the same age range employed in the current study, D1- vs. D2-dopamine receptor expressing MSNs showed differences in passive membrane properties including input resistance, the membrane time constant, and rheobase (Planert et al. 2013). We found no sex differences in passive membrane properties, only in active properties such as the slope of the FI curve and AP threshold and AHP peak amplitude. There were also no signs of bimodality in any property. Thus, the recorded sex differences in MSN intrinsic excitability do not match those reported for rat MSN subtypes and are not likely driven by differential sampling.

A related question is whether both MSN subtypes show sex differences in intrinsic excitability. Future studies should directly test this question. In the meantime, we sorted the present dataset for each sex to find neurons with the 10 lowest and 10 highest values for three attributes that differ between rat MSN subtypes: input resistance, membrane time constant, and rheobase (Planert et al. 2013). We did this to bias each dataset toward containing unequal proportions of MSN subtypes. We then used two-way ANOVAs to analyze whether the FI slope of these neurons significantly differed in both the low value and high value groups of input resistance, membrane time constant, and rheobase (analysis not shown). For all three attributes sex was a significant source of variation. However, there was no inter-

action between sex and attribute amplitude. None of the attributes demonstrated a bimodal distribution. This is consistent with the hypothesis that sex differences may be present in both MSN subtypes. Supporting this, both MSN subtypes express non-nuclear, membrane-associated, estrogen receptor α and β expression in the dorsal striatum (Almey et al. 2012; Grove-Strawser et al. 2010; Kuppens and Beyer 1999; Mermelstein et al. 1996; Schultz et al. 2009; Toran-Allerand et al. 1992). Large area estradiol infusions into female dorsal striatum modulates sensorimotor performance (Becker et al. 1987), paced mating behavior (Xiao and Becker 1997), and learning and memory tasks (Zurkovsky et al. 2007). This indicates that estradiol activation of both MSN subtypes does not appear to compromise estradiol-induced changes in striatal-mediated behaviors. Furthermore, the dorsal striatum as a whole is necessary for aspects of maternal behavior (Henschen et al. 2013; Keer and Stern 1999). We thus tentatively speculate that sex differences in intrinsic excitability are present in both D1- and D2-dopamine receptor expressing MSNs with the full acknowledgement that this prediction needs to be empirically tested.

Changes in intrinsic membrane excitability can occur either independently or in concert with other neuronal attributes such as changes in extracellular nonsynaptic glutamate levels and/or synaptic input (Otaka et al. 2013; Schulz 2006; Suska et al. 2013; Wolf 2010; Zakon 1998). In the current study, no sex differences in mEPSC properties were detected. Possible sex differences in excitatory or inhibitory synaptic input have not yet been examined in adult dorsal striatum or in prepubertal nucleus accumbens. Interestingly, Woolley and colleagues found increased mEPSC frequency and spine density in female rat adult MSNs located in the nucleus accumbens core and to a lesser extent in the nucleus accumbens shell, and that these properties are modulated by cocaine exposure (Forlano and Woolley 2010; Wissman et al. 2012; Wissman et al. 2011). Increased spine density has since been detected in female medium spiny neurons in human nucleus accumbens (Sazdanovic et al. 2013). MSN spine density is modulated by estradiol in adult female hamsters and rats in the nucleus accumbens core but not in other striatal regions (Peterson et al. 2014; Staffend et al. 2011). Given that changes in synaptic input are often linked with changes in intrinsic excitability (Ishikawa et al. 2009; Wolf 2010), this raises the possibility that estradiol or cocaine exposure also differentially modulates intrinsic excitability by sex. While no sex differences in excitatory synaptic input were found by the current study, we do note that both intrinsic excitability and synaptic input may be reorganized during puberty, similar to glutamatergic synaptic input in the medial amygdala (Cooke 2011; Cooke and Woolley 2009), or modulated by adult hormone profile (Woolley 2007; Wu et al. 2011). These questions all represent outstanding avenues of research, in addition to determining ionic mechanism, the possible roles of dopamine and dopamine receptors, and whether the sex differences reported here are generated through a genetic/epigenetic mechanism or via the organizational influences of early steroid sex hormone exposure/absence (McCarthy and Arnold 2011).

ACKNOWLEDGMENTS

We thank Drs. Paul Mermelstein and Robert Meisel and the 2011 Grass Laboratory for support.

GRANTS

NC State University Start-up Funds (J. Meitzen), Grass Foundation Fellowship (J. Meitzen), Howard Hughes Medical Institute Undergraduate Science Education grant (C. A. Hauser, PI: D. Shea).

DISCLOSURES

No conflicts of interest, financial or otherwise, are declared by the author(s).

AUTHOR CONTRIBUTIONS

Author contributions: D.M.D., J.C., J.A.W., and J.M. performed experiments; D.M.D., J.C., J.A.W., C.A.H., and J.M. analyzed data; D.M.D. and J.M. interpreted results of experiments; D.M.D. and J.M. prepared figures; D.M.D., J.C., J.A.W., C.A.H., and J.M. edited and revised manuscript; D.M.D., J.C., J.A.W., C.A.H., and J.M. approved final version of manuscript; J.M. conception and design of research; J.M. drafted manuscript.

REFERENCES

- Almey A, Filardo EJ, Milner TA, Brake WG. Estrogen receptors are found in glia and at extranuclear neuronal sites in the dorsal striatum of female rats: evidence for cholinergic but not dopaminergic colocalization. *Endocrinology* 153: 5373–5383, 2012.
- Arnold AP, Lusis AJ. Understanding the sexome: measuring and reporting sex differences in gene systems. *Endocrinology* 153: 2551–2555, 2012.
- Babayan AH, Kramar EA. Rapid effects of oestrogen on synaptic plasticity: interactions with actin and its signalling proteins. *J Neuroendocrinol* 25: 1163–1172, 2013.
- Baufreton J, Atherton JF, Surmeier DJ, Bevan MD. Enhancement of excitatory synaptic integration by GABAergic inhibition in the subthalamic nucleus. *J Neurosci* 25: 8505–8517, 2005.
- Becker JB. Direct effect of 17 beta-estradiol on striatum: sex differences in dopamine release. *Synapse* 5: 157–164, 1990.
- Becker JB. Hormonal influences on sensorimotor function. In: *Behavioral Endocrinology*, edited by Becker JB, Breedlove SM, Crews D. Cambridge, MA: MIT Press, 2002, p. 497–525.
- Becker JB, Hu M. Sex differences in drug abuse. *Front Neuroendocrinol* 29: 36–47, 2008.
- Becker JB, Perry AN, Westenbroek C. Sex differences in the neural mechanisms mediating addiction: a new synthesis and hypothesis. *Biol Sex Diff* 3: 14, 2013.
- Becker JB, Snyder PJ, Miller MM, Westgate SA, Jenuwine MJ. The influence of estrous cycle and intrastratial estradiol on sensorimotor performance in the female rat. *Pharmacol Biochem Behav* 27: 53–59, 1987.
- Beery AK, Zucker I. Sex bias in neuroscience and biomedical research. *Neurosci Biobehav Rev* 35: 565–572, 2011.
- Belleau ML, Warren RA. Postnatal development of electrophysiological properties of nucleus accumbens neurons. *J Neurophysiol* 84: 2204–2216, 2000.
- Bobzean SA, DeNobrega AK, Perrotti LI. Sex differences in the neurobiology of drug addiction. *Exp Neurol* 259C: 64–74, 2014.
- Breedlove SM, Arnold AP. Sexually dimorphic motor nucleus in the rat lumbar spinal cord: response to adult hormone manipulation, absence in androgen-insensitive rats. *Brain Res* 225: 297–307, 1981.
- Breedlove SM, Hampson E. Sexual differentiation of the brain and behavior. In: *Behavioral Endocrinology*, edited by Becker JB, Breedlove SM, Crews D. Cambridge, MA: MIT Press, 2002, p. 75–114.
- Cahill L. Why sex matters for neuroscience. *Nat Rev Neurosci* 7: 477–484, 2006.
- Calhoun JC. *The Ecology and Sociology of the Norway Rat*. U. S. Department of Health, Education, and Welfare, PHS no. 1008, 1962.
- Campi KL, Jameson CE, Trainor BC. Sexual dimorphism in the brain of the monogamous California mouse (*Peromyscus californicus*). *Brain Behav Evol* 81: 236–249, 2013.
- Carroll ME, Anker JJ. Sex differences and ovarian hormones in animal models of drug dependence. *Horm Behav* 58: 44–56, 2010.
- Chen X, Grisham W, Arnold AP. X chromosome number causes sex differences in gene expression in adult mouse striatum. *Eur J Neurosci* 29: 768–776, 2009.
- Cooke BM. Synaptic reorganization of the medial amygdala during puberty. *J Neuroendocrinol* 23: 65–73, 2011.
- Cooke BM, Woolley CS. Effects of prepubertal gonadectomy on a male-typical behavior and excitatory synaptic transmission in the amygdala. *Dev Neurobiol* 69: 141–152, 2009.
- Cooke BM, Woolley CS. Sexually dimorphic synaptic organization of the medial amygdala. *J Neurosci* 25: 10759–10767, 2005.
- Cosgrove KP, Mazure CM, Staley JK. Evolving knowledge of sex differences in brain structure, function, and chemistry. *Biol Psychiatry* 62: 847–855, 2007.
- Cummings JA, Jagannathan L, Jackson LR, Becker JB. Sex differences in the effects of estradiol in the nucleus accumbens and striatum on the response to cocaine: neurochemistry and behavior. *Drug Alcohol Depend* 135: 22–28, 2014.
- De Vries GJ. Minireview: Sex differences in adult and developing brains: compensation, compensation, compensation. *Endocrinology* 145: 1063–1068, 2004.
- Di Paolo T. Modulation of brain dopamine transmission by sex steroids. *Rev Neurosci* 5: 27–41, 1994.
- Dorris DM, Hauser CA, Minnehan CE, Meitzen J. An aerator for brain slice experiments in individual cell culture plate wells. *J Neurosci Meth* 238: 1–10, 2014.
- Eckel LA, Houtp TA, Geary N. Spontaneous meal patterns in female rats with and without access to running wheels. *Physiol Behav* 70: 397–405, 2000.
- Farries MA, Meitzen J, Perkel DJ. Electrophysiological properties of neurons in the basal ganglia of the domestic chick: conservation and divergence in the evolution of the avian basal ganglia. *J Neurophysiol* 94: 454–467, 2005.
- Farries MA, Perkel DJ. Electrophysiological properties of avian basal ganglia neurons recorded in vitro. *J Neurophysiol* 84: 2502–2513, 2000.
- Farries MA, Perkel DJ. A telencephalic nucleus essential for song learning contains neurons with physiological characteristics of both striatum and globus pallidus. *J Neurosci* 22: 3776–3787, 2002.
- Fattore L, Melis M, Fadda P, Fratta W. Sex differences in addictive disorders. *Front Neuroendocrinol* 35: 272–284, 2014.
- Forlano PM, Woolley CS. Quantitative analysis of pre- and postsynaptic sex differences in the nucleus accumbens. *J Comp Neurol* 518: 1330–1348, 2010.
- Gale SD, Perkel DJ. Physiological properties of zebra finch ventral tegmental area and substantia nigra pars compacta neurons. *J Neurophysiol* 96: 2295–2306, 2006.
- Geller SE, Koch A, Pellettieri B, Carnes M. Inclusion, analysis, and reporting of sex and race/ethnicity in clinical trials: have we made progress? *J Womens Health (Larchmt)* 20: 315–320, 2011.
- Gertler TS, Chan CS, Surmeier DJ. Dichotomous anatomical properties of adult striatal medium spiny neurons. *J Neurosci* 28: 10814–10824, 2008.
- Ghahramani NM, Ngun TC, Chen PY, Tian Y, Krishnan S, Muir S, Rubbi L, Arnold AP, de Vries GJ, Forger NG, Pellegrini M, Vilain E. The effects of perinatal testosterone exposure on the DNA methylome of the mouse brain are late-emerging. *Biol Sex Diff* 5: 8, 2014.
- Giorgi FS, Galanopoulou AS, Moshe SL. Sex dimorphism in seizure-controlling networks. *Neurobiol Dis* 72B: 144–152, 2014.
- Gorski RA, Gordon JH, Shryne JE, Southam AM. Evidence for a morphological sex difference within the medial preoptic area of the rat brain. *Brain Res* 148: 333–346, 1978.
- Grove-Strawser D, Boulware MI, Mermelstein PG. Membrane estrogen receptors activate the metabotropic glutamate receptors mGluR5 and mGluR3 to bidirectionally regulate CREB phosphorylation in female rat striatal neurons. *Neuroscience* 170: 1045–1055, 2010.
- Henschen CW, Palmiter RD, Darvas M. Restoration of dopamine signaling to the dorsal striatum is sufficient for aspects of active maternal behavior in female mice. *Endocrinology* 154: 4316–4327, 2013.
- Hille B. *Ion Channels of Excitable Membranes*. Sunderland, MA: Sinauer, 2001.
- Hosseini-Kamkar N, Morton JB. Sex differences in self-regulation: an evolutionary perspective. *Front Neurosci* 8: 233, 2014.
- Hu M, Watson CJ, Kennedy RT, Becker JB. Estradiol attenuates the K⁺-induced increase in extracellular GABA in rat striatum. *Synapse* 59: 122–124, 2006.

- Huang GZ, Woolley CS. Estradiol acutely suppresses inhibition in the hippocampus through a sex-specific endocannabinoid and mGluR-dependent mechanism. *Neuron* 74: 801–808, 2012.
- Ishikawa M, Mu P, Moyer JT, Wolf JA, Quock RM, Davies NM, Hu XT, Schluter OM, Dong Y. Homeostatic synapse-driven membrane plasticity in nucleus accumbens neurons. *J Neurosci* 29: 5820–5831, 2009.
- Keer SE, Stern JM. Dopamine receptor blockade in the nucleus accumbens inhibits maternal retrieval and licking, but enhances nursing behavior in lactating rats. *Physiol Behav* 67: 659–669, 1999.
- Kole MH, Stuart GJ. Signal processing in the axon initial segment. *Neuron* 73: 235–247, 2012.
- Kourrich S, Thomas MJ. Similar neurons, opposite adaptations: psychostimulant experience differentially alters firing properties in accumbens core versus shell. *J Neurosci* 29: 12275–12283, 2009.
- Kuppers E, Beyer C. Expression of estrogen receptor-alpha and beta mRNA in the developing and adult mouse striatum. *Neurosci Lett* 276: 95–98, 1999.
- Mani SK, Reyna AM, Alejandro MA, Crowley J, Markaverich BM. Disruption of male sexual behavior in rats by tetrahydrofuran diols (THF-diols). *Steroids* 70: 750–754, 2005.
- Markaverich B, Mani S, Alejandro MA, Mitchell A, Markaverich D, Brown T, Velez-Trippe C, Murchison C, O'Malley B, Faith R. A novel endocrine-disrupting agent in corn with mitogenic activity in human breast and prostatic cancer cells. *Environ Health Perspect* 110: 169–177, 2002.
- McCarthy MM, Arnold AP. Reframing sexual differentiation of the brain. *Nat Neurosci* 14: 677–683, 2011.
- Meitzen J, Perry AN, Westenbroek C, Hedges VL, Becker JB, Mermelstein PG. Enhanced striatal β 1-adrenergic receptor expression following hormone loss in adulthood is programmed by both early sexual differentiation and puberty: a study of humans and rats. *Endocrinology* 154: 1820–1831, 2013.
- Meitzen J, Pfelepsen KR, Stern CM, Meisel RL, Mermelstein PG. Measurements of neuron soma size and density in rat dorsal striatum, nucleus accumbens core and nucleus accumbens shell: differences between striatal region and brain hemisphere, but not sex. *Neurosci Lett* 487: 177–181, 2011.
- Meitzen J, Weaver AL, Brenowitz EA, Perkel DJ. Plastic and stable electrophysiological properties of adult avian forebrain song-control neurons across changing breeding conditions. *J Neurosci* 29: 6558–6567, 2009.
- Mermelstein PG, Becker JB, Surmeier DJ. Estradiol reduces calcium currents in rat neostriatal neurons via a membrane receptor. *J Neurosci* 16: 595–604, 1996.
- Mu P, Moyer JT, Ishikawa M, Zhang Y, Panksepp J, Sorg BA, Schluter OM, Dong Y. Exposure to cocaine dynamically regulates the intrinsic membrane excitability of nucleus accumbens neurons. *J Neurosci* 30: 3689–3699, 2010.
- Nottebohm F, Arnold AP. Sexual dimorphism in vocal control areas of the songbird brain. *Science* 194: 211–213, 1976.
- Nunez JL, McCarthy MM. Androgens predispose males to GABAA-mediated excitotoxicity in the developing hippocampus. *Exp Neurol* 210: 699–708, 2008.
- O'Donnell P, Grace AA. Physiological and morphological properties of accumbens core and shell neurons recorded in vitro. *Synapse (New York, NY)* 13: 135–160, 1993.
- Ober C, Loisel DA, Glad Y. Sex-specific genetic architecture of human disease. *Nat Rev Genet* 9: 911–922, 2008.
- Otaka M, Ishikawa M, Lee BR, Liu L, Neumann PA, Cui R, Huang YH, Schluter OM, Dong Y. Exposure to cocaine regulates inhibitory synaptic transmission in the nucleus accumbens. *J Neurosci* 33: 6753–6758, 2013.
- Peterson BM, Mermelstein PG, Meisel RL. Estradiol mediates dendritic spine plasticity in the nucleus accumbens core through activation of mGluR5. *Brain Struct Funct* [Epub ahead of print].
- Planert H, Berger TK, Silberberg G. Membrane properties of striatal direct and indirect pathway neurons in mouse and rat slices and their modulation by dopamine. *PLoS One* 8: e57054, 2013.
- Prendergast BJ, Onishi KG, Zucker I. Female mice liberated for inclusion in neuroscience and biomedical research. *Neurosci Biobehav Rev* 40: 1–5, 2014.
- Remage-Healey L. Frank Beach Award Winner: Steroids as neuromodulators of brain circuits and behavior. *Horm Behav* 66: 552–560, 2014.
- Sato SM, Wissman AM, McCollum AF, Woolley CS. Quantitative mapping of cocaine-induced DeltaFosB expression in the striatum of male and female rats. *PLoS One* 6: e21783, 2011.
- Sazdanovic M, Slobodanka M, Zivanovic-Macuzic I, Jeremic D, Tanaskovic I, Milosavljevic Z, Malikovic A, Ognjanovic N, Sazdanovic P, Jovanovic B, Jovanovic J, Todorovic M, Toseveski J. Sexual dimorphism of medium-sized neurons with spines in human nucleus accumbens. *Arch Biol Sci Belgrade* 65: 1149–1155, 2013.
- Schultz KN, von Esenwein SA, Hu M, Bennett AL, Kennedy RT, Musatov S, Toran-Allerand CD, Kaplitt MG, Young LJ, Becker JB. Viral vector-mediated overexpression of estrogen receptor-alpha in striatum enhances the estradiol-induced motor activity in female rats and estradiol-modulated GABA release. *J Neurosci* 29: 1897–1903, 2009.
- Schulz DJ. Plasticity and stability in neuronal output via changes in intrinsic excitability: it's what's inside that counts. *J Exp Biol* 209: 4821–4827, 2006.
- Shen W, Hamilton SE, Nathanson NM, Surmeier DJ. Cholinergic suppression of KCNQ channel currents enhances excitability of striatal medium spiny neurons. *J Neurosci* 25: 7449–7458, 2005.
- Shen W, Hernandez-Lopez S, Tkatch T, Held JE, Surmeier DJ. Kv1.2-containing K^+ channels regulate subthreshold excitability of striatal medium spiny neurons. *J Neurophysiol* 91: 1337–1349, 2004.
- Srivastava DP, Woolfrey KM, Liu F, Brandon NJ, Penzes P. Estrogen receptor activity modulates synaptic signaling and structure. *J Neurosci* 30: 13454–13460, 2010.
- Staffend NA, Loftus CM, Meisel RL. Estradiol reduces dendritic spine density in the ventral striatum of female Syrian hamsters. *Brain Struct Funct* 215: 187–194, 2011.
- Suska A, Lee BR, Huang YH, Dong Y, Schluter OM. Selective presynaptic enhancement of the prefrontal cortex to nucleus accumbens pathway by cocaine. *Proc Natl Acad Sci USA* 110: 713–718, 2013.
- Tepper JM, Sharpe NA, Koos TZ, Trent F. Postnatal development of the rat neostriatum: electrophysiological, light- and electron-microscopic studies. *Dev Neurosci* 20: 125–145, 1998.
- Toran-Allerand CD, Miranda RC, Hochberg RB, MacLusky NJ. Cellular variations in estrogen receptor mRNA translation in the developing brain: evidence from combined [125I]estrogen autoradiography and non-isotopic in situ hybridization histochemistry. *Brain Res* 576: 25–41, 1992.
- Trabzuni D, Ramasamy A, Inran S, Walker R, Smith C, Weale ME, Hardy J, Ryten M. Widespread sex differences in gene expression and splicing in the adult human brain. *Nat Commun* 4: 2771, 2013.
- Villalon Landeros R, Morisseau C, Yoo HJ, Fu SH, Hammock BD, Trainor BC. Corn cob bedding alters the effects of estrogens on aggressive behavior and reduces estrogen receptor-alpha expression in the brain. *Endocrinology* 153: 949–953, 2012.
- Walker QD, Rooney MB, Wightman RM, Kuhn CM. Dopamine release and uptake are greater in female than male rat striatum as measured by fast cyclic voltammetry. *Neuroscience* 95: 1061–1070, 2000.
- Wissman AM, May RM, Woolley CS. Ultrastructural analysis of sex differences in nucleus accumbens synaptic connectivity. *Brain Struct Funct* 217: 181–190, 2012.
- Wissman AM, McCollum AF, Huang GZ, Nikrodhanond AA, Woolley CS. Sex differences and effects of cocaine on excitatory synapses in the nucleus accumbens. *Neuropharmacology* 61: 217–227, 2011.
- Wolf ME. The Bermuda Triangle of cocaine-induced neuroadaptations. *Trends Neurosci* 33: 391–398, 2010.
- Woodruff TK, Kibbe MR, Paller AS, Turek FW, Woolley CS. Commentary: “Leaning in” to support sex differences in basic science and clinical research. *Endocrinology* 155: 1181–1183, 2014.
- Woolley CS. Acute effects of estrogen on neuronal physiology. *Annu Rev Pharmacol Toxicol* 47: 657–680, 2007.
- Wu WW, Adelman JP, Maylie J. Ovarian hormone deficiency reduces intrinsic excitability and abolishes acute estrogen sensitivity in hippocampal CA1 pyramidal neurons. *J Neurosci* 31: 2638–2648, 2011.
- Xiao L, Becker JB. Effects of estrogen agonists on amphetamine-stimulated striatal dopamine release. *Synapse (New York, NY)* 29: 379–391, 1998.
- Xiao L, Becker JB. Hormonal activation of the striatum and the nucleus accumbens modulates paced mating behavior in the female rat. *Horm Behav* 32: 114–124, 1997.
- Yang CF, Shah NM. Representing sex in the brain, one module at a time. *Neuron* 82: 261–278, 2014.
- Zakon HH. The effects of steroid hormones on electrical activity of excitable cells. *Trends Neurosci* 21: 202–207, 1998.
- Zhang XF, Hu XT, White FJ. Whole-cell plasticity in cocaine withdrawal: reduced sodium currents in nucleus accumbens neurons. *J Neurosci* 18: 488–498, 1998.
- Zurkovsky L, Brown SL, Boyd SE, Fell JA, Korol DL. Estrogen modulates learning in female rats by acting directly at distinct memory systems. *Neuroscience* 144: 26–37, 2007.