

Hydrogen Sulfide: The Third Gasotransmitter in Biology and Medicine

Rui Wang

Abstract

The last two decades have seen one of the greatest excitements and discoveries in science, gasotransmitters in biology and medicine. Leading the trend by nitric oxide and extending the trudge by carbon monoxide, here comes hydrogen sulfide (H_2S) who builds up the momentum as the third gasotransmitter. Being produced by different cells and tissues in our body, H_2S , alone or together with the other two gasotransmitters, regulates an array of physiological processes and plays important roles in the pathogenesis of various diseases from neurodegenerative diseases to diabetes or heart failure, to name a few. As a journal dedicated to serve the emergent and challenging field of H_2S biology and medicine, *Antioxidant and Redox Signaling* assembles the most recent discoveries and most provoking ideas from leading scientists in H_2S fields, which were communicated in the First International Conference of H_2S in Biology and Medicine, and brings them to our readers in two Forum Issues. Through intellectual exchange and intelligent challenge with an open-mind approach, we can reasonably expect that sooner rather than later the exploration of metabolism and function of H_2S will provide solutions for many of the biological mysteries of life and pave way for the arrival of many more gasotransmitters. *Antioxid. Redox Signal.* 12, 1061–1064.

THE STUDIES OF GASEOUS SIGNALING molecules, gasotransmitters, have made great progress over the last several decades. In 1987, nitric oxide (NO) was discovered as the first endogenously produced gaseous signaling molecule. Carbon monoxide (CO) was identified in the middle of the 1990s as the second one. It appears that the appreciation of the physiological importance of hydrogen sulfide (H_2S), the third gasotransmitter (13), emerged on the scientific horizon only yesterday. Yet we have witnessed one of the most fascinating and fastest expanding research evolutions in the recent history of biomedical sciences. We have seen new scientific publications on H_2S research weekly. We have welcomed new research teams joining the H_2S field monthly.

It was timing. To highlight the explosive and mesmerizing development in the H_2S research field and summarize our updated understanding of the biological and medical implications of H_2S , the first international conference on H_2S in Biology and Medicine was held in Shanghai, China from June 26–28, 2009.

It was exciting. About 160 scientists and trainees attended this meeting from 15 countries and regions. The scientific scope covered by six symposia and poster sessions spanning from cell differentiation and apoptosis, inflammation, glucose metabolism, wound healing, neurodegenerative diseases,

cardiac ischemic protection, hypertension, development and reproduction, up to potential human hibernation. The platform provided by this conference allowed researchers to seek interdisciplinary collaboration and translational research effort in H_2S study.

Published in these two Forum Issues of *Antioxidant & Redox Signaling* are examples of the topics and debates that were generated during the conference. The enzymatic basis for endogenous production of H_2S , for example, is case in point. As we know now, H_2S can be produced in a variety of cells, tissues, organs, and systems. Two key enzymes in the transsulfuration pathway, cystathionine β -synthase (CBS) and cystathionine γ -lyase (CSE), have been consistently shown to produce H_2S , pyruvate, and ammonium, using L-cysteine and homocysteine as the substrates (13). In some tissues, both CBS and CSE function to catalyze H_2S production, but in others only one of the two suffices for the production of this gasotransmitter. The existence of additional enzymes in mammalian cells for catalyzing the production of H_2S has been speculated but not substantiated. In this Issue of ARS, Kimura gives a detailed description of 3-mercaptopyruvate sulfurtransferase (3MST). This third enzyme in H_2S production has been identified in brain and vascular endothelium. Using both L-cysteine and alpha-ketoglutarate as substrates, 3MST along

with cysteine aminotransferase (CAT) produces H₂S. As 3MST is mostly localized in neurons but CBS is in astrocytes, these two enzymes play different roles in regulating brain production of H₂S. In blood vessel wall, 3MST can be found in both vascular smooth muscle cells and endothelial cells, whereas CAT is only localized to vascular endothelial cells. As such, it appears that only vascular endothelium cells in blood vessel wall can use 3MST to produce H₂S since the enzymatic function of 3MST requires the co-existence of CAT.

The role of H₂S in the digestive system under physiological or pathophysiological conditions is another interesting topic, especially given the fact that the gastrointestinal (GI) tract contains the highest concentration of H₂S than any other systems in our body. Sulfur-reducing bacteria in the lumen of the large intestine produce H₂S as high as 3 mM through fermentation of sulfur-containing amino acids and intestinal sulfomucin metabolism. GI tissues and cells *per se* also produce H₂S through the action of CBS and/or CSE. As well known, H₂S contributes to the homeostatic control of GI mucosal defence and repair. This protective effect has been ascribed to the anti-inflammatory effects of H₂S, the inhibition of leukocyte adherence to the vascular endothelium, and leukocyte migration to sites of inflammation being examples. If the endogenous production of H₂S from the GI tract is inhibited, as Wallace summarized in his review article in this Issue, the mucosal integrity will be jeopardized and ulcerative and inflammatory diseases of the GI tract would occur (10). While acknowledging some well-studied protective effects of H₂S on the GI tract, another review article by Linden *et al.* in this issue summarizes the current challenges we are facing (8). The regulation of GI production of H₂S is questioned. The target molecules or signaling pathways of H₂S in the GI tract have not been clear. The interaction of bacteria-yielded H₂S and H₂S from the cells of the GI tract and its impact on the GI tract function should be examined. Within the context of the GI tract investigation, Cao *et al.* report in this Issue, for the first time, the expression of both CSE and CBS as well as the endogenous production of H₂S in colon cancer cells and colonic tissues (3). Looking for the endogenous regulation of GI production of H₂S? Butyrate is one of these regulators. Presented in the mammalian colon at high millimolar concentrations, butyrate serves as an energy substrate for colonic epithelial cells, where it is metabolized by mitochondrial β -oxidation. It has been shown that this short chain fatty acid in the GI tract inhibits cell growth and differentiation of several human colon cancer cell lines. Butyrate increases cellular production of H₂S by upregulating the expression of CBS (3). Thus yielded H₂S also reduces the viability of colon cancer cells. Whereas both butyrate and H₂S share the same inhibitory effect on cell growth, these two endogenous substances regulate cancer cell growth through different mechanisms.

Also communicated in this Issue is altered H₂S metabolism in different diseases, including insulin resistance syndrome, hyperhomocysteinemia (HHcy)-induced cardiomyocyte injury, and regional ischemia damages. The report by Chang *et al.* shed light on the interaction of H₂S with methylglyoxal (MG) in vascular smooth muscle cells (VSMCs) (4). Being a metabolite of sugar, protein, and fatty acid, MG is formed in virtually all mammalian cells. Chang *et al.* found that cellular production of H₂S is decreased by MG in a dose-dependent manner in VSMCs. This effect is correlated to the inhibition of CSE by MG. On the other hand, NaHS (a H₂S donor) lowers

cellular MG level and decreases MG-induced reactive oxygen species production. Since MG plays a pathogenic role in the development of insulin resistance syndrome, including hypertension, diabetes, and obesity, the involvement of H₂S in these very diseases may be modulated, at least partially, by MG and vice versa. A direct molecule-to-molecule reaction between MG and H₂S would change the structure and function of both involved molecules. It would be interesting to map out the *in vivo* conditions for this reaction to occur and the new molecular structure yielded. The effect of H₂S on HHcy-induced cardiomyocyte damage is reported by Wei *et al.* (14). This study correlated the plasma homocysteine level and endogenous H₂S production with endoplasmic reticulum (ER) stress in cardiomyocytes. The methionine overload-induced HHcy leads to two events in the tested rats, a significant increase in ER stress and a decreased endogenous production of H₂S. Once exogenous H₂S is supplied to these tested animals, cardiomyocytes are rescued from ER stress. Increased ER stress is also observed after endogenous H₂S production is inhibited. Since ER stress is not a patent to HHcy, this study predicts a protective role of H₂S in many other pathological situations in which ER stress is elevated. Whether and how H₂S protects regional ischemia-damaged tissues and organs are examined by Wang M-J *et al.* in this Issue (12). These authors used an *in vivo* hind limb ischemia rat model with unilateral femoral artery occluded for 4 weeks. NaHS supplementation to these animals leads to significant collateral vessel growth, increased capillary density, and improved regional tissue blood flow in ischemic hind limb muscles in comparison with the controls. Increased expression of endothelial growth factor (VEGF) and VEGF receptor 2 and phosphorylation of Akt in ischemic hind limb muscle were also resulted from NaHS treatment. As such, a pro-angiogenic effect of H₂S is indicated. This finding may have significant impact on the management of various ischemic damages to different tissues and organs.

The cytoprotective effects of H₂S against hypoxic and inflammation injuries and the underlying signaling mechanisms are the focus of the next Forum Issue of ARS. Calvert *et al.* present a detailed review on recent discoveries on H₂S-elicited amelioration of *in vitro* or *in vivo* myocardial and hepatic ischemia-reperfusion injury (2). This anti-ischemic protection can be achieved by both applying exogenous H₂S and upregulating the endogenous production of H₂S. In line with this notion, invention of new vehicles to promote endogenous H₂S production can find its important cytoprotective applications. The effects of three novel cysteine containing compounds against rat acute myocardial ischemia are reported in the next Forum Issue (11). These compounds preserve the activities of antioxidant enzymes and reduce the formation of the lipid peroxidation product in ventricular tissues. These beneficial effects are likely mediated by the CSE-catalyzed H₂S production as the inhibition of CSE abolished the protective effects of these three compounds.

In addition to the antioxidant protection, the oxygen-sensing ability of endogenous H₂S would be another mechanism for the cytoprotective effects of this gasotransmitter. The hypothesis that H₂S may be an oxygen-sensing molecule (9), if approved, would fill up a major knowledge gap in our understanding on how the oxygen-sensing cells sense changes in oxygen level and then launch counter reactions to regain a desirable oxygen level. Chemoreceptors and chromaffin cells

are known oxygen-sensing cells in vertebrate cardiorespiratory systems. It is speculated that oxidation of endogenously produced H₂S triggers the cardiorespiratory reflexes and relaxes smooth muscles to increase local blood perfusion and/or ventilation (4). Though this theory is still in the hypothetical stage, lines of supportive evidence have been amounted. Not only hypoxia and H₂S have similar effects on selective types of cells, increased H₂S production augments hypoxic responses. Moreover, H₂S has been shown to be rapidly consumed by O₂ in the range of intracellular/mitochondrial PO₂. Based on these findings, we may need to re-evaluate the regulation of oxygen utilization and mitochondrial function and to add H₂S in the signaling chain for energy production and usage inside the cells.

The challenge facing H₂S researchers can also be well represented by the controversial observations on the role of H₂S in inflammation. In the next Issue of ARS Forum, two articles are specifically devoted to this topic. Bhatia reviews the interaction of H₂S with different inflammation mediators, especially Substance P (1). Based on the observations from his own team and the others on different animal inflammation models, including sepsis and acute pancreatitis, Bhatia believes that H₂S plays a pro-inflammatory role linked to the metabolism of Substance P. On the other hand, the anti-inflammatory effect of H₂S has been reported by many research teams. One explanation for the seemingly opposite roles of H₂S in inflammation resides in the concentration of H₂S being tested and the organs where H₂S is produced as well. Whiteman *et al.* approach this issue from a unique angle (15). These authors examined the effect of a novel H₂S donor (GYY4137) on lipopolysaccharide (LPS)-mediated inflammation. The slow releasing of H₂S from GYY4137 inhibited LPS-induced release of pro-inflammatory mediators and increased the synthesis of the anti-inflammatory chemokine IL-10. The rapid releasing of H₂S from a sulfide salt NaHS at high concentrations, in contrast, increased the synthesis of pro-inflammatory factors. It is concluded that the rate of H₂S generation is important, if not critical, for determining whether H₂S is a pro- or anti-inflammation substance.

As the knowledge base of the metabolism and function of H₂S in our body has been enlarged rapidly over the last decade, more molecular mechanisms underlying the cellular effects of H₂S have been unmasked. K_{ATP} channel is the earliest identified signaling molecular target of H₂S (18). The stimulation of K_{ATP} channels by H₂S has been introduced as the molecular basis for H₂S-elicited cardiac protection (2), blood pressure lowering (17), or insulin release inhibition (16). The mechanism by which H₂S changes K_{ATP} channel function, however, has been a puzzle. Jiang *et al.* have combined the whole-cell patch-clamp technique with a mutagenesis approach to investigate this puzzle (5). They for the first time discovered that H₂S increases K_{ATP} channel currents by specifically interacting with the SUR subunit of K_{ATP} channel complex. Furthermore, they demonstrated that selective cysteine residues located on the extracellular loop of SUR subunit were chemically modified by H₂S. With these observations, the stimulation of K_{ATP} channels by H₂S now can be understood as the consequence of altered configuration of K_{ATP} channel complex. In addition to the stimulation of K_{ATP} channels, H₂S also acts on other ion channels in different types of cells. Li *et al.* report that H₂S inhibits big-conductance calcium-activated K⁺ (BK_{Ca}) channels in type I glomus cells of

the carotid body (7). It was found that both hypoxia and NaHS decreased the whole-cell BK_{Ca} channel currents significantly and reversibly. Inhibition of CBS activity in type I glomus cells abolished the sensitivity of BK_{Ca} channels to hypoxia. The possibility is thus raised that endogenous H₂S generated by CBS activity may mediate hypoxic modulation of BK_{Ca} channels in chemoreceptor cells.

Exploration of biological importance of H₂S is one of the best examples of how curiosity drives research and innovation. If our fellow researchers and readers become even more curious about H₂S, the purpose of the First International Conference on H₂S in Biology and Medicine and these two related Forum Issues of ARS will have been well served. The unsolved puzzles about the metabolism and functions of H₂S are still many. As food for thought, I will mention some of the puzzles here. The enzymatic reactions leading to H₂S production have not been fully understood. The emergence of 3MST is greatly welcomed but the functional importance of this enzyme in the endogenous production of H₂S remains to be determined. Whether 3MST can produce H₂S under physiological conditions is uncertain, given that its full activity requires an alkaline condition (pH 9.7) (6). Inflammation can be promoted by H₂S and suppressed by the same gasotransmitter. Can we find way to suppress its pro-inflammatory role and enhance its anti-inflammatory role? Increased tissue content of H₂S protects heart from ischemia/reperfusion damage. Meanwhile, high concentration of H₂S has high potential to disrupt electron transport chain in cardiac mitochondrion. One of the intriguing and promising applications of H₂S for cardiac protection is to use it as a preconditioning as well as a postconditioning agent. To score a practical point, a postconditioning appears more feasible than the preconditioning application of H₂S but the effectiveness comparison needs to be conducted. As soon as an oxygen-sensing role is given to H₂S, the questions are immediately asked as why hypoxia and H₂S share the same function if H₂S is a protective factor against hypoxic damage. H₂S releasing can be affected by hypoxia, and so does the function of H₂S. Does H₂S itself have the same or opposite biological affects with different oxidation levels? Or in another word does H₂S lose its cellular effects in the presence of oxygen? Then, let me post the question in everyone's mind—what would be the dividing line between the toxic and physiological levels of H₂S?

The leadership of Dr. Yizhun Zhu and his local organization committee at the Faculty of Pharmacy, Fudan University, China, are key to the success of the First International Conference of H₂S in Biology and Medicine. Dr. Philip Moore from King's College (UK) and Dr. Hideo Kimura from National Institute of Neuroscience (Japan) offered their invaluable support in helping me edit these two Issues of ARS Forum.

Research on H₂S in biology and medicine is entering its most exciting period. The only way to maintain the momentum and ultimately unmask the importance of H₂S to our health to its fullest is to let curiosity take the lead and let the inspiration to enquire endure. The farewell message on the Whole Earth Catalogs (1974, edition) should be the welcome message in H₂S study: "Stay hungry, stay foolish".

References

1. Bhatia M. Hydrogen sulfide and Substance P in inflammation. *Antioxid Redox Signal* 12: 1191–1202, 2010.

2. Calvert JW, Coetzee WA, and Lefer DJ. Novel insights into hydrogen sulfide-mediated cytoprotection. *Antioxid Redox Signal* 12: 1203–1217, 2010.
3. Cao Q, Zhang L, Yang G, Xu C, and Wang R. Butyrate-stimulated H₂S production in colon cancer cells. *Antioxid Redox Signal* 12: 1101–1109, 2010.
4. Chang T, Untereiner A, Liu J, and Wu L. Interactions of methylglyoxal and hydrogen sulfide in rat vascular smooth muscle cells. *Antioxid Redox Signal* 12: 1093–1100, 2010.
5. Jiang B, Tang G, Cao K, Wu L, and Wang R. Molecular mechanism for H₂S-induced activation of K_{ATP} channels. *Antioxid Redox Signal* 12: 1167–1178, 2010.
6. Kimura H. Hydrogen sulfide: From brain to gut. *Antioxid Redox Signal* 12: 1111–1123, 2010.
7. Li Q, Sun B, Wang X, Jin Z, Zhou Y, Dong L, Jiang L-H, and Rong W. A crucial role of hydrogen sulfide in oxygen sensing *via* modulating large conductance calcium-activated potassium channels. *Antioxid Redox Signal* 12: 1179–1189, 2010.
8. Linden DR, Levitt MD, Farrugia G, and Szurszewski JH. Endogenous production of H₂S in the gastrointestinal tract: Still in search of a physiological function. *Antioxid Redox Signal* 12: 1135–1146, 2010.
9. Olson KR and Whitfield NL. Hydrogen sulfide and oxygen sensing in the cardiovascular system. *Antioxid Redox Signal* 12: 1219–1234, 2010.
10. Wallace JL. Physiological and pathophysiological roles of hydrogen sulfide in the gastrointestinal tract. *Antioxid Redox Signal* 12: 1125–1133, 2010.
11. Wang Q, Wang X-L, Liu H-R, Rose P, and Zhu Y-Z. Protective effects of cysteine analogues on acute myocardial ischemia: Novel modulators of endogenous H₂S production. *Antioxid Redox Signal* 12: 1155–1165, 2010.
12. Wang M-J, Cai W-J, Li N, Ding Y-J, Chen Y, and Zhu Y-C. The hydrogen sulfide donor NaHS promotes angiogenesis in a rat model of hind limb ischemia. *Antioxid Redox Signal* 12: 1065–1077, 2010.
13. Wang R. Two's company, three's a crowd: Can H₂S be the third endogenous gaseous transmitter? *FASEB J* 16: 1792–1798, 2002.
14. Wei H, Zhang R, Jin H, Liu D, Tang X, Tang C, and Du J. Hydrogen sulfide attenuates hyperhomocysteinemia-induced cardiomyocytic endoplasmic reticulum stress in rats. *Antioxid Redox Signal* 12: 1079–1091, 2010.
15. Whiteman M, Li L, Rose P, Tan C-H, Parkinson DB, and Moore PK. The effect of hydrogen sulfide donors on lipopolysaccharide-induced formation of inflammatory mediators in macrophages. *Antioxid Redox Signal* 12: 1147–1154, 2010.
16. Wu L, Yang W, Jia X, Yang G, Duridanova D, Cao K, and Wang R. Pancreatic islet overproduction of H₂S and suppressed insulin release in Zucker diabetic rats. *Lab Invest* 89: 59–67, 2009.
17. Yang G, Wu L, Jiang B, Yang W, Qi J, Cao K, Meng Q, Mustafa AK, Mu W, Zhang S, Snyder SH, and Wang R. H₂S as a physiologic vasorelaxant: Hypertension in mice with deletion of cystathionine gamma-lyase. *Science* 322: 587–590, 2008.
18. Zhao W, Zhang J, Lu Y and Wang R. The vasorelaxant effect of H₂S as a novel endogenous gaseous K_{ATP} channel opener. *EMBO J* 20: 6008–6016, 2001.

Address correspondence to:

Dr. Rui Wang
Office of the Vice-President (Research)
Lakehead University
955 Oliver Road
Thunder Bay, Ontario P7B 5E1
Canada

E-mail: rwang@lakeheadu.ca

Date of first submission to ARS Central, October 7, 2009; date of final revised submission, October 20, 2009; date of acceptance, October 21, 2009.

Abbreviations Used

3MST = 3-mercaptopyruvate sulfurtransferase
BK_{Ca} channel = big-conductance calcium-activated K⁺ channel
CAT = cysteine aminotransferase
CBS = cystathionine beta-synthase
CSE = cystathionine gamma-lyase
CO = carbon monoxide
ER stress = endoplasmic reticulum stress
GI tract = gastrointestinal tract
HHCy = hyperhomocysteinemia
H₂S = hydrogen sulfide
K_{ATP} channel = ATP-sensitive K⁺ channel
MG = methylglyoxal
NO = nitric oxide
VEGF = endothelial growth factor
VSMCs = vascular smooth muscle cells